

Winter Cereal Rye Cover Crop Effect on Cash Crop Yield

Year 8

Iowa Learning Farms and Practical Farmers of Iowa



Summary

Farmers reported that in 55 of 59 site-years, properly managed cover crops had little to no negative effect on corn and soybean yield (and actually increased soybean yield in 7 site-years and corn yield in 2 site-years).

Cooperators

Bill Buman, Harlan
Randy Caviness, Greenfield
Jim Funcke, Jefferson
Devan Green, Conrad
Rick Juchems, Plainfield
Rob Davis & Darwin Pierce,
Whiterock Conservancy, Coon Rapids
Mark Pokorny, Clutier
George Schaefer, Kalona
Jerry Sindt, Holstein
Rob Stout, West Chester
Gary & Dave Nelson, Fort Dodge
Kelly Tobin, New Market

Project Timeline

2008-2016 (8th year report)

This project was funded by the State Soil Conservation Committee, the Iowa Department of Agriculture and Land Stewardship and the Leopold Center for Sustainable Agriculture. Additional outreach and education funding came from a NCR-SARE grant, Walton Family Foundation and the Iowa Learning Farms.

Methods

- 12 sites over the course of this study with six participating in 2016. All sites are in corn-soybean rotations.
- Cooperators establish and maintain replicated strips the length of their field for duration of the study. Each replication has one strip with cover crops and one without cover crops.
- Cooperators seed cereal rye cover crop in the fall of 2015 aerially or with a drill at seeding rates ranging between 56-60 lb/ac.
- Cover crop termination was accomplished with herbicide applied prior to cash crop planting the following spring.

Table 1. Farm location, cover crop management, and cash crop for the 2016 growing season.

Location	Cover Crop planting date	Cover crop planting method	Cover crop seeding rate	Cover crop termination method	Cash crop	Planting date
Plainfield (NE Iowa)	10/8/15	Drilled	56 lb/ac	Herbicide	Corn	5/8/16
Coon Rapids (West central Iowa)	10/20/15	Drilled	60 lb/ac	Herbicide	Corn	4/25/16
Holstein (NW Iowa)	9/9/15	Aerial	60 lb/ac	Herbicide	Soy	5/18/16
West Chester (SE Iowa)	11/4/15	Drilled	56 lb/ac	Herbicide	Corn	4/25/16
New Market (SW Iowa)	10/14/15	Drilled	56 lb/ac	Herbicide	Corn	4/22/16
Jefferson (West Central Iowa)	11/1/15	Drilled	60 lb/ac	Herbicide	Soy	5/15/16

Results

Cover crop biomass

Above-ground cover crop biomass was determined at most locations at the time of cover crop termination (Table 2). Over the years, aboveground cover crop biomass at locations ranged from trace amounts to 2,407 lb/ac prior to corn and from 55- 2,475 lb/ac prior to soybeans. Cover crop was typically terminated 7-10 days prior to planting.


Location	Cover crop biomass (lb/ac)	Sampling date
Plainfield	680.12	4/29/16
Coon Rapids	515.42	4/19/16
Holstein	834.71	5/12/16
West Chester	454.24	4/14/16
New Market	231.14	4/19/16
Jefferson	398.90	4/27/16

Table 2. Mean cover crop aboveground biomass samples prior to termination in 2016.

Corn yields 2016

In general, corn yields were equivalent regardless of cover crop treatment as determined by statistical analysis (t-test). Only at Jefferson (2009), Coon Rapids (2010) and Harlan (2010) were corn yields reduced in the cover crop strips. It should be noted that these instances occurred only in the first two growing seasons of the trial. Cooperators identified insufficient cover crop termination (Jefferson '09) or improper planter settings (Coon Rapids, Harlan '10) as reasons for the average yield decrease of 27 bu/ac. In the remaining cases, corn yields were mostly not affected by the cover crop (Figure 1). In 2016, corn yields were statistically improved by 3 and 19 bu/ac at West Chester and New Market, respectively.


Figure 1. Trends with respect to cover crop effect on corn yields at 10 site-years from 2009 to 2010 and 24 site-years from 2011 to 2016.



Soybean yields 2016

Soybean yields were typically equivalent regardless of cover crop treatment as determined by statistical analysis (t-test). In seven cases, however, soybean yields were improved by the cover crop. Increase in soybean yield ranged from 3 to 8 bu/ac with an average increase of 5 bu/ac in these cases. As with corn, soybean yield was also mostly not affected by the cover crop (Figure 2). Only at Coon Rapids (2013) were soybean yields reduced in the cover crop strips. The cooperator identified planter setting as a reason for the yield difference. Soybean yields in 2016 ranged between 63 and 64 bu/ac at two locations.

Figure 2. Trends with respect to cover crop effect on soybean yields at 6 site-years from 2009 to 2010 and 19 site-years from 2011 to 2016.



Cover crop effect on cash crop yield trends

Since 2008, there have been 34 site-years dedicated to determining the effect on corn yields and 25 site-years to determine the effect on soybean yields. After their first year of introducing cereal rye into their operations, the farmer partners made adjustments to their planter settings to handle more residue and planned to terminate the cover crop 10-14 days before planting to minimize negative impacts on yield. After eight years in the study, the farmer partners have reported mostly no effect of the cereal rye cover crop on corn and soybean yield.

For more detailed information on the project and Year 5 report, see "Winter Cereal Rye Cover Crop Effect on Cash Crop Yield – Year 5" on these websites:

ILF: <https://www.iowalearningfarms.org/content/cover-crop-research>

PFI: <http://practicalfarmers.org/farmer-knowledge/research-reports/2014/winter-cereal-rye-cover-crop-effect-cash-crop-yield/>



1201 Sukup Hall
Ames, Iowa 50011-3080
Liz Juchems: 515-294-5429
ejuchems@iastate.edu
www.iowalearningfarms.org



600 Fifth Street
Ames, Iowa 50010-6071
Stefan Gailans: 515-232-5661
stefan@practicalfarmers.org
www.practicalfarmers.org